Bull, Basrah nat, Hist, Mus, Vol, 1 No, 1: 1974 June

TURSIOPS ADUNCUS BOTTLE NOSED DOLPHIN: A NEW RECORD FOR ARAB GULF; WITH NOTES ON CETACEA OF THE REGION.

L

Khalaf Al -- Robaae

Director, Basrah Natural History Museum

One dolphin was collected on 15 January 1974 from Ras Al — Mataf, Arab Gulf by a fishing boat. Mr: Bashir Zahroon of the Iraqi National Fisheries Company has presented the specimen to the Museum:

The dead dolphin was black on the back and light gray on the belly. The black around the eye extended anteriorly (see Plate 1). The specimen has been identified as a male Tursiops aduncus (Ehrenburg, 1832). The measurements are given below:

Total length 2 m.

Pectoral fin 35 cm.


Height of dorsal fin 21 cm.


Width of the flukes 54 cm.

The lower jaw contained 26 teeth on each side while the upper jaw had only 25 teeth on each side; each tooth measured about one centimeter in diameter:


The present collection of T . aduncus from the Arab Gulf forms the first record of this species from this region . The distribution of this species as given by (Hershkovitz 1966) is as follows:


Platei





Platei





Distribution.

Indian Ocean: from the Rea Sea and South Africa to Bay of Bengal and Australia Pacific: from New Zealand, Australia, Indonesia, and China Seas, on the west to American waters from Baja California to Chile, on the east; Atlantic Ocean; from Gulf of San Matias, Rio Negro, Argentina to Rio de la Plata and in the Rio Urguway to Paysandu, Urguway, and off Rio Grande do Sul, Brazil:

The Fauna of Cetacea From Arabian Gulf

Several species of Whales and Dolphins have been recorded from the Arabian Gulf by different authors .

The latest information available to me is given below.

The Dolphins are localy known by the name "Dughoos" in Arabic and the Whales as "Hoodt".

1 — Sotalia lentigenosa (Owen 1866) Steno (Souse) lentigenosa , Gray 1866 ; proc zool . Soc . London , 1866 .

A male Dolphin was caught in a branch of Khor - Al - Zubair (Arab Gulf -- Iraqi Waters) by fisherman on 25 July 1967, is the second record of this species. The specimen (length 220 cm) has been preserved in the Mammalia Section in the Natural History Museum in Basrah.

Recorded by (Al -- Robaae, 1970).

2 — Sotalia plumbea -- (G: Cuvier, 1856).

Delphinus plumbeus G. Cuvier 1826 Regeanima, ed. 2, 1.

228 ftn. Pucheran 1959 Rev. Mag. Zool (2), 8:145, 449.

(Characters) Bruyns 1960, Malayan Nat. Journ. 14:159

Fig. pp. 164 — 5 (animal) (Indian -- Arabian Sea, Arabian Gulf, Gulf of Aden, Red Sea, Suez Canal). Local name

"Dorfeel" (Kuwait).

One speciment was recorded by Buryns in 1960 from the Kuwait Coast (Hershkovitz, 1966).

Recorded by (Buryns, 1960).

3 — Sotalia fergosoni Lydeker 1903 .

Lydeker 1903, Journ., Bombay Nat. Hist. Soc., 15: 411. Plate D (Animal).

A female was caught by fishermen on the coast of Shauikh Secondary School -- Kuwait (Arab Gulf) in spring 1962 (length 178 cm., teeth 36/33): The animal is preserved in the Audio -- visual aids Department -- Kuwait. The animal was described and figured as a new record from Arabian Gulf by the Author and his personal communication with Dr. P. C. Fraser (British Museum -- Natural History).

Recorded by (A) - Robaae) .

4 — Stenella malayana (Lesson 1826 a) . Delphinus maray--

Voyage autour du monde sur la Corvotte -- La Coquille, Zoology, I (1): 184 Atlas, pl. 9.

Seen by the author in the Fao area, swimming with the dolphin Sotollia lentigenosa. It is gray in colour.

Distinguished from S. lentigenosa by its colour and its dorsal fin. The animal was described and figured as new record for Arabian Gulf by the author.

Recorded by (Al -- Robaae) .

5 — Pseudorea crassidens (Owen 1846).

False killer whale, El -- Hoodt El -- Kattil (Arab).

Pseudorea crassidens -- Owen 1846.

A history of British fossil mammals and birds, P. 518, Fig. 213 (Skull).

One specimen was found in 1964 in Dowha at 40 Km . South of Kuwait (Arab -- Gulf -- Kuwaiti Waters) in muddy area . The skeleton of this specimen is displayed in the Natural History Museum -- Kuwait . The skeleton was mounted in Audio -- Visual Department -- Kuwait .

6 — Balaenoptera musculus -- (Linnaeus 1758).

Great Blue Whale, El -- Hoodt El -- Azrak (Arabic).

(Balana) musculus Linnaeus 1758 . Syst . Nat . ed . 10 .

1:76 . True 1898, Proc. U.S. Nat. Mus., 21629.

Type History: -- Thoma 1911 Proc. Zool. Soc. London,

1911, 2: 156 (Type History).

A dead whale was found entangled in the submarine telegraph cable, and for years a large whale haunted the harbour of Masket in Arabia. Another whale was found in "Kadguma" on the Kuwaiti Coast on June 6, 1963 in muddy area. Its skeleton was mounted by the Natural History Museum in Shuaikh Secondary School, Kuwait.

Recorded by . (Blanford) dead specimen .

(Al -- Robaae, 1969) an skeleton.

7 — Balaenoptera edeni -- Anderson (Bryde's Whale).

Balaemoptera edeni -- Anderson . Anatomical and Zoological Research comprising an account of zoological results of two edpeditions to Western Greece 1868 and 1875 . London , pp . 551--564 .

One whale has been identified as male. Bryde's whale which was stranded on the Island infront of Um Qasr (Arab Gulf -- Iraqi Waters).

Recorded by : (N . Mahdi , 1967) .

(Al -- Robaae , 1969) .

8 — Megaptera novaeangliae (Borowski 1781).

Humpback Whale: El -- Hoodt El -- Ahdab (Arabic)

Ba'aena novaeangliae -- Borowski , Gemmein . Naturgesch , des Tierreichs Berlin 2 , 1 -- (1781) 21 .

A vertebra and a rib of a whale that are said to have been found about 1954 during the construction of a wall near Fao are now in the Iraqi Natural History Museum (Bashir Allous in litt). There is an old report that a Turkish gunboat killed this whale about a century ago in the Shatt - Al - Arab.

Recorded by : (R . Hatt , 1959) :

, in the increase of a

SUMMARY

The first record of Tursiops aduncus (Ehrenberg, 1832) from Arab Gulf is reported together with the status of the Cetacean Fauna of this region.

يتناول هــــذا البحث تسجيل نوع من الدولفينات (عنق الزجاج) للمرة الاولى من الخليج العربي وكذلك يشمل مسحا أوليا لجميع الحيتان والدولفينات المسجلة لحد الآن في الخليج العربي .

ACKNOWLEDGEMENT

I am grateful to Dr. P. J. H. van Bree, Curator of Mammals, University of Amsterdam for helping me in the identification of T. aduncus and to Mr. Bashir Zaharron of the Iraqi National Fisheries Company for presenting the above specimen to the Museum.

BIBLIOGRAPHY

- 1 Al -- Robaae . K . (1969) Brye's whale on the coast of Iraq Z . F . Saugetier Kunde . Bd . 34 (1969) H . 2 . st: 120 125 .
- - 3 False Killer whale "pseudorca crassidens"

- A new record for Arab Gulf . Iraq Nat : Hist : Mus :

- 6 Bruyns . (1960) Malayan Nat . Journ . 14 : 159 . figs pp . 164 -- 5 (animal (Indian) .
- 7 Rice, D. W. and V. B. Scheffer 1968. A List of the Marine Mammals of the World. U. S. Fish and Wildlife Service Special Scientific Report Fisheries No. 579 Washington.
- 8 Hatt.R. (1959). The Mammalia of Iraq. Misc Publ.

 Mus. Zool. Univ. Michigan. Ann Arbor 106:1-33.
- 9 Hershkovitz, ph. (1966) Catalog of Living Whales:
 Bull: 246. Smithsonian Institute United States National Museum.
 Washington, D. C.
- 10 Mahdi . N . (1967); First record of Bryde's whale Balaenoptera edeni Anderson from Arab with notes on earlier literature . Bull . Iraq Nat . Hist . Mus . 3 (7) . 1-6 .
- 11 Norris . K. S. (-- 966) Whales , Dolphins , and Porpoises . University of California Press . Berkelep and Los Angles .
- 21 Mahdi and P. V. George (1969) A systematic list of the vertebrate of Iraq. Bull. Iraqi Nat Mus. publication No. 26 · pp · 1 104:
- 13 Walker . E. R. and associntes (1964) . Mammalia of the world . Vol . II . John Hopkins . Baltimore 647 -- 1500 .